

## review

# Gambling Disorder in the College Student-Athlete Population: An Overview

Donald E. Nowak, Jr.<sup>1</sup>

<sup>1</sup>Department of Counseling and Clinical Psychology, Medaille College, Buffalo, NY, USA

### Abstract

This review examines research from 1991 to the present regarding college student-athlete gambling addiction and disorder issues, with an emphasis on prevalence rates, motivations, and comorbid disorders, as well as National Collegiate Athletic Association (NCAA) national studies and derivative research. Subsets of the college student-athlete population, specifically minority athletes, are also examined. Databases PsycINFO, PsycARTICLES, ERIC, SPORTDiscus, MEDLINE, and Dissertation Abstracts International (ProQuest), were searched for possible contributions to this review. It was determined that student-athletes, and male student-athletes in particular, are vulnerable to disordered gambling problems, which, if university administration and athletic departments do not address, may result in severe negative consequences for the student-athlete. The research suggests that, for the most part, student-athletes have a higher rate of pathological gambling than non-athletes, though the rate of “normal” gambling behavior is about the same. Additionally, it appears that athletes in certain high profile team sports (football, basketball, etc.), as well as athletes belonging to a minority group, are more likely to report problems with gambling than their counterparts. Recommendations for working with student-athletes with a gambling disorder, as well as directions for future research in this burgeoning area, are offered. These proposals include screening for the disorder by mental health professionals and counsellors, as well as training for coaches and financial aid personnel.

**Keywords:** gambling disorder, college student-athletes, prevalence, addiction, NCAA

### Résumé

Cette étude fait l'examen de la recherche effectuée de 1991 à nos jours sur les problèmes de dépendance et de désordre chez les étudiants-athlètes, en mettant l'accent sur les taux de prévalence, les motivations et les troubles concomitants, ainsi

que les études nationales de la National Collegiate Athletic Association et des travaux de recherche dérivés. Des sous-ensembles de la population d'étudiants-athlètes universitaires, en particulier des athlètes faisant partie de minorités, sont également soumis à l'étude. Des recherches ont été faites dans les bases de données PsycINFO, PsycARTICLES, ERIC, SPORTDiscus, MEDLINE et Dissertation Abstracts International (ProQuest) pour trouver d'éventuelles contributions à la présente étude. On a établi que les étudiants-athlètes, masculins en particulier, sont vulnérables aux problèmes de jeu compulsifs, et s'ils ne sont pas pris en main par l'administration universitaire et les départements sportifs, ces troubles peuvent avoir de graves conséquences pour eux. La recherche laisse entendre que, pour la plupart, les étudiants-athlètes ont un taux de jeu pathologique plus élevé que les non-athlètes, bien que le taux de jeu « normal » soit à peu près le même. De plus, il semble que les athlètes de certains sports d'équipe de haut niveau (football, basketball, etc.), ainsi que les athlètes appartenant à un groupe minoritaire, sont plus susceptibles de montrer des problèmes de jeu que leurs homologues. Des recommandations sont faites pour travailler avec des étudiants-athlètes ayant un trouble du jeu, ainsi que des orientations pour de futures recherches dans ce domaine en progression. Ces propositions comprennent le dépistage du trouble par des professionnels de la santé mentale et des conseillers, ainsi que la formation des entraîneurs et du personnel de l'aide financière.

---

### **Introduction**

The economic downturn and volatility of the past decade have together provoked an unprecedented number of financially struggling state and national governments to turn to and rely more heavily upon, gambling. The expansion of lotteries, the increasing approval of more casinos, and the establishment of racetrack slot machines have all occurred as a means to generate these earnings. (Stuart, 2011). In addition to these governmental agencies, academic institutions and religious organizations have also turned to gambling to meet increasing expenditures amid the backdrop of imposed cutbacks (Iancu, Lowengrub, Dembinsky, Kotler, & Dannon, 2008). Even those countries that do not officially permit gambling for its citizens because of cultural or religious reasons often sanction gambling venues for foreign visitors (e.g., Malaysia and South Korea) (Hodgins, Stea, & Grant, 2011). This rapid growth in gambling is not only driving up profits for gambling organizers, but also is producing a dramatic increase in gambling disorder (Hodgins et al., 2011). The concomitant cost to society is staggering: one Baylor University researcher estimated that addicted gamblers cost the United States alone between \$32.4 billion and \$53.8 billion per year (Stuart, 2011).

Many adolescents begin gambling at an age earlier than they do other risky behaviors, such as smoking and alcohol use (Ladouceur, Dube, & Bujold, 1994). Thus, by the time they reach their college years, many young men and women have

already begun gambling, whereas other young persons may simply become involved in gambling as a “rite of passage” while in college (Stinchfield, Hanson, & Olson, 2006). Regardless, young adult populations between the ages of 18–25 have become especially vulnerable to gambling problems, with significantly higher rates of participation than those of the general adult population, and as such are often specifically targeted by advertising campaigns (Annenberg Public Policy Center, 2005; Lesieur et al., 1991; Nowak & Aloe, 2013; Shaffer, Donato, LaBrie, Kidman, & LaPlante, 2005).

The rise in problem gambling has been most notably reflected in the increase in college counseling centers reporting cases of students facing issues such as insurmountable debt, depression, and academic problems, as well as college student-athletes in jeopardy of expulsion for illegal activities related to gambling (Martin, Nelson, & Galucci, 2016; National Collegiate Athletic Association, 2003; Oster & Knapp, 1998). The problem of gambling disorders can prove especially noteworthy among college and university students, many of whom have the resources, proximity, free time, and general desire to become involved in the myriad options of gambling such as casinos, Internet gambling websites, daily fantasy sports (DFS) online, poker games on- and off-campus, sports gambling, instant scratch-off tickets, and state lotteries. This situation does not begin to include the vast number of illegal and informal modes of gambling (often involving a bookmaker) that may expose a student to personal safety issues, as well as to the obvious and inevitable monetary losses.

College students are particularly susceptible to developing gambling disorder issues because of the confluence of several different factors. Those factors create a “perfect storm,” composed of what this researcher terms “The Five A’s”: (1) *age*, with the college years being associated with a wide range of risky behaviors (LaBrie, Shaffer, LaPlante, & Weschler, 2003); (2) *availability* of wide-scale legal (and illegal) gambling, including online gambling; (3) *acceptability* of gambling operated by various government entities and integrated into mainstream culture; (4) *advertising* and media which promote, glorify, and glamorize gambling as a legitimate sport; and (5) *access* to monetary funds, especially from student loans and through numerous credit card solicitations. The result is a population group specifically targeted by both the media and advertisers, which is especially vulnerable to gambling problems.

### **College Student-Athletes and Gambling**

The area of pathological gambling in college student-athletes was generally ignored in favor of an initial focus on treatment of pathological gambling in adults and on addiction and prevention in adolescents. But gambling scandals in the late 1990s at Arizona State University, Boston College, and Northwestern University caught the attention of the National Collegiate Athletic Association (NCAA), its member institutions, the press, and fans of intercollegiate athletics. Other researchers began to believe that college students might well represent the segment of our population with the highest rate of pathological gambling (Weiss & Loubier, 2008).

## Method

The process involved in obtaining information about problem gambling and probable pathological gambling and its prevalence among college and university student-athletes was thorough and exhaustive. To identify all possible studies from 1987 to the present, the researcher used the search terms “gambling,” “college students,” and “student-athletes” as the primary search terms, as well other synonyms, such as “gaming” for “gambling”; “disordered” and “compulsive” for “pathological”; and “university students” for “college students.” The following online databases were closely examined using the search terms outlined above: PsycINFO, PsycARTICLES, ERIC, SPORTDiscus, and MEDLINE. Dissertation Abstracts International (ProQuest) was also searched for possible contributions, and did in fact suggest several possible studies for inclusion. Bibliographies and references from the past three published syntheses in this area of interest (Blinn-Pike, Worthy, & Jonkman, 2007; Nowak & Aloe, 2013; Shaffer, Hall, & Vander Bilt, 1999) were examined as well, for possible additions to the set of studies. All three syntheses assisted in the review.

Additionally, this researcher used the terms noted above in search functions on the online library of the Responsible Gambling Council ([www.responsiblegambling.org](http://www.responsiblegambling.org)), as well as on the online gambling library of the International Gambling Research Institute ([www.gamlib.org](http://www.gamlib.org)). Both digital libraries provide comprehensive resources for scholarly articles in the gambling research field, an area of inquiry which is still growing, and is relatively nascent in terms of generating academic work, specifically in the realm of college students’ gambling behavior.

## Studies

One of the earliest published articles looking at the gambling behaviors and attitudes towards risk-taking in general with student-athletes was by Cross, Basten, Hendrick, Kristofic, and Schaffer (1998) at the University of Michigan. The researchers surveyed 648 NCAA Division I football and men’s basketball players, with the two goals of (1) better understanding student-athlete gambling, and (2) encouraging “more extensive research that may lead to preventive measures in the future” (p. 432). The results suggested that 25.5% of the college student-athletes gambled on intercollegiate athletic events—in itself is an NCAA violation carrying with it a one-year suspension from athletic participation (Cullen & Latessa, 1996)—and 3.7% had wagered on a sporting event in which they had participated, an NCAA violation with a mandatory lifetime ban.

Additionally, Cross et al. found that the men’s basketball and football players who gambled on sports had significantly different attitudes towards risk-taking than those student-athlete counterparts who had not participated in gambling activities: much more permissive attitudes in general. This level of permissiveness towards risk-taking was higher in the football players than the basketball players, supporting one hypothesis of the researchers: the more physical nature of the sport of football

predicts such a result, because the players were, presumably, high-sensation seeking individuals to begin with.

Another study that looked specifically at the group of college student-athletes (Kerber, 2005) not only attempted to ascertain levels of problem and pathological gambling, but also attitudes towards gambling and the modes of gambling they preferred, as well as certain of the predictors of gambling problems. Using the South Oaks Gambling Screen (SOGS) and the Gambling Attitude Scale (GAS), Kerber surveyed 636 student-athletes at three Midwestern universities, and found that while nearly one-quarter (24.1%) of the college athletes claimed never to have gambled, the observed SOGS scores (i.e., those scores  $\geq 3$ ) indicated that 15% exhibited either problem or pathological gambling.

The SOGS was originally intended to screen for pathological gambling in clinical settings, but over the past quarter-century, has since expanded to other purposes, populations, and settings, including prevalence estimate studies of pathological gambling in the general population (Stinchfield, 2002). The past-year self-report version has indicated good overall classification accuracy (.96), with better sensitivity (.99) than specificity (.75), suggesting in turn that the SOGS tends to more often identify false positives (Stinchfield, 2002), a common limitation mentioned by researchers.

Using a multiple regression analysis to predict this total SOGS score from six variable sets, it was concluded that the variables that best predict gambling problems (via a SOGS score) were: (1) frequency of gambling behavior; (2) number of family members or friends with gambling problems; (3) race (i.e., being a minority group member); and (4) age (older rather than younger). No correlation could be found between grade point average and SOGS scores, but student-athletes who were in a fraternity or sorority were found to have higher rates of pathological and problem gambling. The most frequent modes of gambling, for those who did in fact gamble, were in games of skill, such as betting on golf, or playing cards (specifically poker) for money.

**NCAA national study and derivative research.** As stated earlier, the NCAA had taken a pointed interest in the gambling behaviors of college student-athletes, particularly because of the damage that could be done to its reputation and those of its member institutions. To that end, the NCAA commissioned the 2003 NCAA National Study on Collegiate Sports Wagering and Associated Health-Risk Behaviors, a self-administered, voluntary, and anonymous survey, which was returned by 20,739 student-athletes. The study comprised 102 questions and was the most comprehensive and first truly national assessment of college student-athletes ever undertaken (Huang, Jacobs, Derevensky, Gupta, & Paskus, 2007a). Four of the published research articles that used this valuable information are examined in this literature review.

Huang et al. (2007), in the first of a series of articles using this extensive NCAA data set, attempted to examine prevalence rates of problem and pathological gambling,

as well as the most popular forms of gambling for student-athletes, and the particular NCAA sports which were most susceptible to having gambling problems and issues. The researchers found that past-year prevalence was consistently higher among male student-athletes than it was among their female counterparts, compatible with all gambling research. On the basis of DSM-IV Gambling Screen method, 4.3% of men, and 0.4% of women were identified as problem or pathological gamblers.

Athletes involved in golf, ice hockey, and lacrosse were seen to have the highest rates of participants who reported wagering on any sporting event. In addition, student-athletes in gender-specific sports wagered more than did their counterparts in uni-sex sports. As found previously by Kerber (2005), the three most popular forms of gambling were card playing, games of skill, and lotteries. Of particular interest to the NCAA, a small number (1.1%) of student-athletes were asked to directly influence the outcome of a sporting event because of a sports wagering debt. However, Huang et al. (2007b) caution the reader in the brief limitations section that it is quite reasonable to assume that certain of these numbers are underreported “because of the sensitive nature of the questions asked, especially with athletic and scholarship eligibility at stake” (p. 98).

Ellenbogen, Jacobs, Derevensky, Gupta, and Paskus (2008) used the 2003 NCAA survey data to determine whether certain student-athletes were more prone to frequent or problem gambling behavior. Examining gender, race, type of sport played, and gambling mode, among many correlates, Ellenbogen et al. found that Hispanic males reported the highest problem and pathological rates and that the percentage of gamblers was highest among Division III student-athletes, followed by Divisions II and I, respectively. In addition, members of team sports were more likely to gamble than student-athletes in individual sports. Student-athletes in high profile sports were more likely than other student-athletes simply to gamble, to gamble weekly, be at-risk gamblers, be pathological gamblers, and to place more money on sports wagers.

In attempting to explain certain of these findings, the researchers echoed the findings of Cross et al. in stating that it is plausible that high-profile sports attract individuals who are particularly competitive and risk-takers, and that these personality types are generally associated with problem gambling. Ellenbogen et al. found, in both this study and in a previous one, that minorities may be especially vulnerable to gambling problems (Ellenbogen, Gupta, & Derevensky, 2007); the authors explain, however, that the risk to college athletes in high-profile sports goes beyond greater representation of ethnic minorities. High profile sports in which minorities are less represented (e.g., ice hockey, golf) display comparable gambling rates. They reasoned that “at least part of the reason for the high prevalence of gambling problems lies in the nature of high-profile sports, the personalities of the athletes attracted to these sports, or a combination of both” (p. 359).

Huang, Jacobs, Derevensky, Gupta, and Paskus (2007b) returned to the same NCAA data set in examining the connection between gambling and health risks

among college student-athletes. The researchers reiterated certain of the findings from their earlier study, but also considered such health risk behaviors such as alcohol, cigarette, and marijuana usage, along with other drug use, eating disorders, and incidents of unprotected sex. Students were variously classified as either non-gambler, social gambler, problem gambler, or pathological gambler.

The results indicated a general upward trend in the data that suggested that as the level of gambling-related problems increased, so did the prevalence of substance use and abuse, gorging and vomiting, and risky sexual practices with multiple partners. Cross-group comparisons by gambler type (as outlined above) were all significant. These findings led additional credence to the idea of risk-taking and permissiveness towards risk being a vital predictor of possible problem gambling in college student-athletes. Huang et al. (2007b) concluded by suggesting the need for “multi-faceted initiatives to tackle these risk behaviors simultaneously” (p. 397).

One specific study looked at the particular issue of heavy episodic drinking (HED) using the NCAA data in relation to DSM-based problem gambling (Huang, Jacobs, & Derevensky, 2011). The study aimed to “empirically examine the prevalence patterns and odds of at-least-weekly alcohol use and HED in relation to various levels of gambling severity in college student-athletes” (p. 302). Different studies had suggested the link between gambling and drinking, especially in college students, and particularly with student-athletes, but none had examined them empirically, especially with such a large sampling (almost 21,000 respondents).

Similar to other studies, and not unexpectedly, the researchers found that males had a higher prevalence of gambling and rates of drinking alcohol than females. Univariate and multivariate logistic regression models revealed that problem gambling was the strongest covariate of at-least-weekly HED. The prevalence of alcohol use increased significantly as gambling level severity increased. Additionally, the steep increase in relative risk also suggested a possible quadratic relationship between gambling level and HED. The researchers concluded by urging health care providers, college administrators, and athletics personnel to develop evidence-based policies and initiatives to curb college drinking and gambling problems, and to incorporate gambling as a risk factor in future investigations of college drinking.

One of the main limitations mentioned or at least alluded to in most of the previously mentioned student-athlete studies is the generalizability of the findings to the population of college students in general. However, several studies have been published which have comparatively examined the gambling behaviors of non-athlete college students and student-athletes. Three of these studies are reviewed here, with certain surprisingly disparate results.

**Comparative studies.** Engwall, Hunter, & Steinberg (2004) surveyed 1,350 undergraduates at the four campuses of Connecticut State University in the fall of 2000, using a modified South Oaks Gambling Screen. The researchers determined that 18% of the men and 4% of the women had had at least three negative life

consequences because of gambling. These negative consequences included feeling guilty about gambling, participating more than they had intended, and using money earmarked for other expenses (rent, car, food) for gambling instead. Similar Huang et al.'s (2011) findings above, Engwall et al. (2004) determined that those participants were identified as problem gamblers were also significantly likely to be heavy drinkers, report negative consequences of drinking activity, and be regular cigarette and marijuana users.

Problem gambling was also related to binge eating and greater use of weight-control efforts. Engwall et al. also found that the percentage of male team athletes (for this research, students were asked if they participated in intercollegiate or club sports) involved in problem and pathological gambling (26%,  $N = 122$ ) to be significantly higher than the rate among non-athletes. This same pattern was also indicated in female athletes when compared to their non-athlete counterparts. Male and female student-athletes alike also gambled more frequently on card games, in sports betting, and in games of skill, as had been reported in certain of the other studies cited in this literature review.

Another study to look at gambling and other high risk behaviors in college students (Stuhldreher, Stuhldreher, & Forrest, 2007) surveyed over 1,000 Pennsylvanian college students, with part of the research devoted to noting patterns of gambling among student-athletes as compared to non-athletes. The researchers found that significantly more athletes (17%) than non-athletes (9%) reported ever gambling ( $p < .01$ ), and also had more gambling debt (5%) than did non-athletes (1%;  $p < .001$ ). However, a significantly higher percentage of athletes actually sought help for gambling problems compared with non-athletes (7% vs. 4%;  $p < .05$ ). The researchers also found that these significant differences were gender-specific to the men in the sample only. Not surprisingly, the number of females reporting gambling problems in the sample was so few as to render no significant differences.

Stuhldreher et al. (2007) also raised a question which had not been mentioned in any of the reviewed articles in this paper, namely, "Should the measure (of gambling problems) be lifetime prevalence, past-year prevalence, or prevalence during school?" (p. 79). This question is an important one that should be addressed in most research on this subject of gambling with student-athletes and non-athletes alike, because including lifetime, or even summertime into a student's assessment may incorporate different responses based on the students' past behaviors that researchers may or may not want to be included in the scope of such studies.

One research article that did not find any real statistical difference between college student-athletes and a student cohort. Weinstock, Whelan, Meyers, & Watson (2007) attempted to replicate previous prevalence work on student-athlete gambling, as well as examine risk factors for gambling behavior and pathology. The researchers stated that their study improved on previous studies by strictly defining student-athletes as intercollegiate athletes, assessing gambling at four universities from geographically diverse areas, and using a comparison cohort of non-athlete students.



A total of 736 student-athletes in 15 NCAA sports and a cohort of 1,071 non-athlete students from the same universities participated in the study.

The results proved somewhat surprising on more than one level. First, no significant differences were determined between student-athletes and non-athletes in terms of gambling frequency, use of a bookmaker, and disordered (i.e., problem and pathological) gambling. The only difference from earlier studies was that student-athletes proved actually less likely to engage in sports betting than the student cohort. The most alarming similarity was the lifetime prevalence rate of disordered gambling, with 12% of males and approximately 4% of females identified as disordered gamblers. These figures were considerably higher than any of the other studies reviewed had reported finding, and prompted Weinstock et al. to comment that “the notion that gambling is a university wide phenomenon in which student-athletes require supplementary attention because of the potential harm to intercollegiate athletics” (p. 21).

In the only meta-analysis of its type to date, Nowak (2014) examined 124 independent data estimates retrieved from 72 studies conducted between 1987 and 2015, surveying 41,989 university students and student-athletes worldwide. The estimated proportion of probable pathological gamblers among students was computed at 6.13%, with a 6.46% rate among student-athletes; this difference was not statistically significant. Rates of problem gambling were computed at 10.23% and 8.97%, for students and student-athletes respectively, and in this case were statistically significant. Nowak also found that Black and Hispanic student-athletes were at an increased risk for exhibiting the indicators of problem gambling.

### **Conclusions and Recommendations**

It appears that gambling behavior on university campuses is a problem that does not yield any signs of abating anytime soon. Student-athletes in general, and male student-athletes in particular, are vulnerable to disordered gambling problems, which, if not addressed by university administration and athletic departments, can result in severe negative consequences for the student-athlete, and possibly both the institution and their personal reputations. The prevalence studies referenced here display disparate results in regards to athletes versus students in terms of gambling severity, with most inquiries exhibiting few to no significant differences. One possible reason for this could be underreporting by student-athletes because of the perceived NCAA ramifications as previously noted. For those stakeholders in college athletics, the fact that these rates are just as high in student-athletes as in students (the highest percentage population of gambling disorder) should stand as a cause for concern and attention. Most notably, it appears that athletes in certain high-profile, revenue-generating team sports (football, basketball, etc.) are more likely to report problems with gambling than their counterparts participating in less visible athletic programs.

**College faculty and staff.** It would seem logical that faculty could also benefit from in-service training related to gambling disorders and how to recognize certain

of the signs of a gambling problem, such as lateness or missed classes, declining grades and performance, tiredness, and irritability. While these symptoms could in fact represent a litany of other issues, the main goal is to help faculty be cognizant of the possibility of a gambling problem as a potential cause of such symptoms. University personnel, such as financial aid counsellors, should also be trained in detecting and screening for excessive gambling, as should residence hall directors and assistants, who see and often interact with students in a much different milieu than do faculty and administrators. Those university employees involved in the delivery of health services should also be trained in screening students for mental health problems, including those problems pertaining to gambling, when presenting for physical exams or problems.

**Coaches.** Because of the particular risk to student-athletes, as well as the inherent dangers of damaging an institution's reputation because of gambling-related scandals, college coaches and other members of athletic departments involved in recruiting, training, and coaching students should be provided with basic education on the popularity of sports wagering and the risks associated with gambling. These persons should also be made aware of the signs and symptoms of disordered gambling in the same type of training that other university faculty and staff should be strongly encouraged to participate in. This prescription is particularly strong for those coaches who work in the high-profile NCAA Division I sports on which most gambling activity in Las Vegas and online casinos is focused.

### **Limitations and future directions**

Putting aside the relative dearth of research in the area of college student-athletes and gambling disorder, limitations include, but are not limited to, the previously mentioned possibility of underreporting by student-athletes, as well as missing data in a number of the articles reviewed, as researchers indicated students did not in all cases answer fully what was asked of them. One limitation that kept cropping up was the use of lifetime gambling measures (SOGS; DSM-IV) which could have resulted in a certain amount of distorted information rather than asking specifically about gambling experiences while enrolled in college. Future studies could be better served by inquiring about the college experience exclusively, as well as comparing and contrasting the percentage rates of gambling disorder as per the DSM-5 (APA, 2013) versus the long-standing pathological and problem gambling rates which have been reported on prior to the new classification of this serious disorder which affects many college student-athletes in the United States. By addressing these issues, the larger body of work regarding college students and gambling can be better served not only to understand the scope of the problem, but also how best to address it and by what means, as students' proclivity and access to the myriad options of both legal and illegal forms of gambling are still emerging. In fact, as states seriously consider legalizing sports betting, these gambling opportunities for young people already deeply invested in sports will grow as well.

## References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual* (5th edition). Washington, DC: Author.
- Annenberg Public Policy Center (2005). *The National Annenberg Survey of Youth*. Philadelphia, PA: The University of Pennsylvania.
- Blinn-Pike, L., Worthy, S., & Jonkman, J. N. (2007). Disordered gambling among college students: A meta-analytic synthesis. *Journal of Gambling Studies*, *23*, 175–183. doi:10.1007/s10899-006-9036-2
- Cross, M. E., Basten, J., Hendrick, E. M., Kristofic, B., & Schaffer, E. J. (1998). Student-athletes and gambling: An analysis of attitudes towards risk-taking. *Journal of Gambling Studies*, *14*, 431–439. doi:10.1023/A:1023081228054
- Cullen, F. T., & Latessa, E. J. (1996). *Assessing the extent and sources of NCAA rule infractions: A national self-report study of student-athletes*. Indianapolis, IN: The National Collegiate Athletic Association. doi: 10.1111/j.1745-9133.2012.00840.x
- Ellenbogen, S., Gupta, R., & Derevensky, J. L. (2007). A cross-cultural study of gambling behaviour among adolescents. *Journal of Gambling Studies*, *23*, 25–39. doi:10.1007/s10899-006-9044-2. Retrieved from: <http://youthgambling.mcgill.ca/en/PDF/Publications/2007/A%20Cross-Cultural%20Study%20of%20Gambling%20Behavior%20Among%20Adolescents%202007.pdf>
- Ellenbogen, S., Jacobs, D., Derevensky, J., Gupta, R., & Paskus, T. (2008). Gambling behavior among college student-athletes. *Journal of Applied Sport Psychology*, *20*, 349–362. doi:10.1080/10413200802056685
- Engwall, D., Hunter, R., & Steinberg, M. (2004). Gambling and other risk behaviors on university campuses. *Journal of American College Health*, *52*, 245–255. doi:10.3200/JACH.52.6.245-256
- Hodgins, D. C., Stea, J. N., & Grant, J. E. (2011). Gambling disorders. *Lancet*, *378*, 1874–1884. doi:10.1016/S0140-6736(10)62185-X
- Huang, J. H., Jacobs, D. F., & Derevensky, J. L. (2011). DSM-based problem gambling: Increasing the odds of heavy drinking in a national sample of U.S. college athletes? *Journal of Psychiatric Research*, *45*, 302–308. doi:10.1016/j.jpsychires.2010.07.001
- Huang, J.-H., Jacobs, D. F., Derevensky, J. L., Gupta, R., & Paskus, T. S. (2007a). Gambling and health risk behaviors among U.S. college student-athletes: Findings from a national study. *Journal of Adolescent Health*, *40*, 390–397. doi:10.1016/j.jadohealth.2006.11.146

- Huang, J.-H., Jacobs, D. F., Derevensky, J. L., Gupta, R., & Paskus, T. S. (2007b). A national study on gambling among US college student-athletes. *Journal of American College Health, 56*, 93–99. doi:10.3200/JACH.56.2.93-100
- Iancu, I., Lowengrub, K., Dembinsky, Y., Kotler, M., & Dannon, P. N. (2008). Pathological gambling: An update on neuropathophysiology and pharmacotherapy. *CNS Drugs, 22*, 123–138. Retrieved from: [https://www.researchgate.net/publication/5657504\\_Pathological\\_gambling\\_an\\_update\\_on\\_neuropathophysiology\\_and\\_pharmacotherapy](https://www.researchgate.net/publication/5657504_Pathological_gambling_an_update_on_neuropathophysiology_and_pharmacotherapy)
- Kerber, C. (2005). Problem and pathological gambling among college athletes. *Annals of Clinical Psychiatry, 17*, 243–247. doi:10.1080/10401230500295388
- LaBrie, R. A., Shaffer, H. J., LaPlante, D. A., & Wechsler, H. (2003). Correlates of college student gambling in the United States. *Journal of American College Health, 52*, 53–62. doi:10.1080/07448480309595725
- Ladouceur, R., Dubé, D., & Bujold, A. (1994). Prevalence of pathological gambling and related problems among college students in the Quebec metropolitan area. *The Canadian Journal of Psychiatry/La Revue Canadienne de Psychiatrie, 39*, 289–93.
- Lesieur, H. R., Cross, J., Frank, M., Welch, M., White, C. M., Rubenstein, G., ... Mark, M. (1991). Gambling and pathological gambling among university students. *Addictive Behaviors, 16*, 517–527. doi:10.1016/0306-4603(91)90059-Q
- Martin, R. J., Nelson, S. E., & Gallucci, A. R. (2016). Game on: Past year gambling, gambling-related problems, and fantasy sports gambling among college athletes and non-athletes. *Journal of Gambling Studies, 32*, 567–579. doi:10.1007/s10899-015-9561-y
- National Collegiate Athletic Association. (2003). *NCAA national study on collegiate sports wagering and associated health-risk behaviors*. Indianapolis, IN: Author.
- Nowak, D. E. (2014). *A meta-analytical synthesis and examination of pathological and problem gambling rates among college students and student-athletes* (Doctoral dissertation). Buffalo, NY: State University of New York at Buffalo.
- Nowak, D. E., & Aloe, A. M. (2013). The prevalence of pathological gambling among college students: A meta-analytic synthesis, 2005–2013. *Journal of Gambling Studies, 30*, 1–25. doi:10.1007/s10899-013-9399-0
- Oster, S. L., & Knapp, T. J. (1998). Sports betting among college students: Who bets and how often? *College Student Journal, 32*, 289–292.
- Shaffer, H. J., Donato, A. N., LaBrie, R. A., Kidman, R. C., & LaPlante, D. A. (2005). The epidemiology of college alcohol and gambling policies. *Harm Reduction*

*Journal*, 2, 1–20. doi:10.1186/1477-7517-2-1. Retrieved from: [https://www.researchgate.net/publication/8030017\\_The\\_epidemiology\\_of\\_college\\_alcohol\\_and\\_gambling\\_policies](https://www.researchgate.net/publication/8030017_The_epidemiology_of_college_alcohol_and_gambling_policies)

Shaffer, H. J., Hall, M. N., & Vander Bilt, J. (1999). Estimating the prevalence of disordered gambling behavior in the United States and Canada: A research synthesis. *American Journal of Public Health*, 89, 1369–1376. doi:10.2105/AJPH.89.9.1369. Retrieved from: <https://pdfs.semanticscholar.org/d89c/b6c0213ba15921938d19875d1459d87b9848.pdf>

Stinchfield, R. (2002). Reliability, validity, and classification accuracy of the South Oaks Gambling Screen (SOGS). *Addictive Behaviors*, 27, 1–19. doi:10.1016/S0306-4603(00)00158-1

Stinchfield, R., Hanson, W. E., & Olson, D. H. (2006). Problem and pathological gambling among college students. In McClellan, G. S., Hardy, T. W., & Caswell, J. (Eds.), *Gambling on campus: New directions for student services* (pp. 63–72). San Francisco, CA: Jossey-Bass.

Stuart, E. (2011, August 9). Gambling on the rise: Is America becoming addicted? *Deseret News*. Retrieved from: <http://www.deseretnews.com/article/700169522/Gambling-on-the-rise-Is-America-becoming-addicted.html?pg=all>

Stuhldreher, W. L., Stuhldreher, T. J., & Forrest, K. Z. (2007). Gambling as an emerging health problem on campus. *Journal of American College Health*, 56, 75–83. doi:10.3200/JACH.56.1.75-88

Weinstock, J., Whelan, J. P., Meyers, A. W., & Watson, J. M. (2007). Gambling behavior of student-athletes and a student cohort: What are the odds? *Journal of Gambling Studies*, 23, 13–24. doi:10.1007/s10899-006-9043-3

Weiss, S. M., & Loubier, S. L. (2008). Gambling behaviors of former athletes: The delayed competitive effect. *UNLV Gaming Research and Review Journal*, 12, 53–60. Retrieved from: <https://pdfs.semanticscholar.org/65e2/4d5432f8d3612ea314c61f67ed8635e807bd.pdf>

\*\*\*\*\*

Submitted May 20, 2017; accepted May 24, 2018. This article was peer reviewed. All URLs were available at the time of submission.

For correspondence: Donald E. Nowak, Jr., PhD, CRC, Department of Counseling and Clinical Psychology, Medaille College, Buffalo, New York, USA, 14214. E-mail: [Donald.E.Nowak@medaille.edu](mailto:Donald.E.Nowak@medaille.edu)

Competing interests: None declared.

Ethics approval: Not required.